

# Introduction to Deep Learning for Speech and Language Processing

## Exercise Sheet 2: Math for Machine Learning

28th October 2025

### Notation

We try our best to be consistent with the mathematical notation throughout the course. Here are the most important conventions we use:

Symbol	Example	Description
greek letter	$\alpha$	scalar
lower-case letter	$x$	vector (column vector)
... with one subscript	$x_i$	$i$ -th entry of vector $x$ (scalar)
... with two subscripts	$x_{ij}$	entry in the $i$ -th row, $j$ -th column of a matrix (scalar)
upper-case letter	$M$	matrix
	$M^{-1}$	inverse matrix of matrix $M$
	$v^\top, M^\top$	transposition of a vector or matrix
	$I_n$	$n \times n$ identity matrix
.	$x \cdot y$	dot product of two vectors
superscript	$W^l, b^l$	weight matrix and bias at layer $l$ of a neural network
$\partial$	$\frac{\partial y}{\partial x}$	partial derivative of $y$ with respect to $x$

For derivatives we use the numerator layout:

$$\frac{\partial y}{\partial x} = \begin{bmatrix} \frac{\partial y_1}{\partial x_1} & \frac{\partial y_1}{\partial x_2} & \dots & \frac{\partial y_1}{\partial x_n} \\ \frac{\partial y_2}{\partial x_1} & \frac{\partial y_2}{\partial x_2} & \dots & \frac{\partial y_2}{\partial x_n} \\ \frac{\partial y_m}{\partial x_1} & \frac{\partial y_m}{\partial x_2} & \dots & \frac{\partial y_m}{\partial x_n} \end{bmatrix} \text{ for } x \in \mathbb{R}^n, y \in \mathbb{R}^m$$

### Linear Algebra

#### Linear Independence

##### Exercise 1.

Given are the vectors  $u = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$ ,  $v = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$  and  $w = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$

- (1) Are the vectors  $u$  and  $v$  linearly independent?
- (2) Is the set of all three vectors  $u, v, w$  linearly independent?

## Determinants and Eigenvalues

### Exercise 2.

Given the matrix  $A = \begin{bmatrix} 2 & 4 \\ 0 & 3 \end{bmatrix}$  Compute.

- (1)  $A - \lambda I_2$
- (2)  $\det(A - \lambda I_2)$
- (3) Both eigenvalues of  $A$

## Derivatives

### Exercise 3.

Compute the derivative w.r.t.  $x \in \mathbb{R}$  for the following functions:

- (1)  $f(x) = (-x - 7)^2$
- (2)  $g(x, y) = -8x^2 + 5x + 7y$
- (3)  $h(x) = \begin{bmatrix} 2x^2 \ln(6x) \\ 3x \end{bmatrix}$

### Exercise 4.

Compute the derivative w.r.t.  $x = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} \in \mathbb{R}^2$  for the following functions:

- (1)  $f(x) = 9x_1 + 10x_2$
- (2)  $g(x_1, x_2) = \begin{bmatrix} e^{-4x_1+5} \\ \frac{1}{7}e^{7x_2} \\ 4x_1 9x_2 \end{bmatrix}$

### Exercise 5.

Compute the derivative w.r.t.  $x \in \mathbb{R}^n$  for the following functions:

- (1)  $f(x) = x$